

Jaewon Chang, DO Jared Smith, MD 300 Hoffman Street Elmira, NY 14845 (607) 734-4110

Post Operative Instructions: Total Hip Arthroplasty

You are recovering from reconstructive hip surgery. The following information is to help make your recovery as smooth and rapid as possible.

- 1. Diet resume your normal diet at home. A well balanced diet promotes healing.
- 2. Activity Weight bearing as tolerated on the operative leg. Use your walker as instructed. Continue the strengthening exercises as instructed by your physical therapist.
- 3. Restrictions Total hip precautions remain in effect for 12 weeks after surgery. You must keep a pillow between your knees when in bed. You may not flex your hip greater than 90 degrees and do not cross your legs.
- 4. Incision Leave the incision open to air and the surgical superglue and mesh dressing will peel off over the next 2 weeks. We will remove the rest of it at your 2 week follow up appointment. You may apply a dressing if drainage occurs. If a different dressing was used we will give you instructions at time of discharge.
- 5. Hygiene You may shower but avoid tub bath or hot tub for the first three weeks following surgery. Wash your hands frequently and try to avoid touching your incision.
- 6. Work You may not return to work until cleared by your provider
- 7. You will continue on a blood thinner for four weeks after discharge. We will use Prilosec once daily for GI protection while on aspirin therapy unless you are already taking a similar medication.
- 8. Continue to use the incentive spirometer every 2 hours while awake.
- 9. Constipation is common following surgery and can be prevented by increasing water and fluid intake, increasing fiber in the diet, and by adding Docusate sodium 2 tabs daily while taking narcotic pain medication. You may try warming equal parts of ginger ale and prune juice in the



Sports Medicine

Jaewon Chang, DO Jared Smith, MD 300 Hoffman Street Elmira, NY 14845 (607) 734-4110

microwave for 1-2 minutes. This may provide relief of symptoms. If this is not effective, you may take Milk of Magnesia or over the counter laxatives.

- 10. Follow-up in the Orthopedic office in about 10- 14 days.
- 11. All dental appointments will require oral antibiotics after total joint replacement. You may not have any dental work done for the first 3 months following surgery. We can order this medication for the appointment. Make your dentist aware of your joint replacement.